

Envirocheck[®] LANDMARK INFORMATION GROUP[®]

Artificial Ground and Landslip

Artificial ground is a term used by BGS for those areas where the ground surface has been significantly modified by human activity. Information about previously developed ground is especially important, as it is often associated with potentially contaminated material, unpredictable engineering conditions and unstable ground.

Artificial ground includes:

- Made ground - man-made deposits such as embankments and spoil heaps on the natural ground surface.


- Worked ground - areas where the ground has been cut away such as quarries and road cuttings.

- Infilled ground - areas where the ground has been cut away then wholly or partially backfilled.

- Landscaped ground - areas where the surface has been reshaped.

- Disturbed ground - areas of ill-defined shallow or near surface mineral workings where it is impracticable to map made and worked ground separately.

Mass movement (landslip) deposits on BGS geological maps are primarily superficial deposits that have moved down slope under gravity to form landslips. These affect bedrock, other superficial deposits and artificial ground. The dataset also includes foundered strata, where the ground has collapsed due to subsidence.



Order Details

 Order Number:
 145523120_1_1

 Customer Ref:
 10/1004

 National Grid Reference:
 335080, 389170

 Slice:
 A

 Site Area (Ha):
 0.21


 Search Buffer (m):
 1000

Site Details

Site at, Liverpool, Liverpool



Tel: Fax: Web:



LANDMARK INFORMATION GROUP®


Superficial Geology

BGS 1:10,000 Superficial Deposits are the youngest geological deposits formed during the most recent period of geological time, which extends back about 1.8 million years from the present.

They rest on older deposits or rocks referred to as Bedrock. This dataset contains Superficial deposits that are of natural origin and 'in place'. Other superficial strata may be held in the Mass Movement dataset where they have been moved, or in the Artificial Ground dataset where they are of man-made origin.

Most of these Superficial deposits are unconsolidated sediments such as gravel, sand, silt and clay, and onshore they form relatively thin, often discontinuous patches or larger spreads.

Superficial Geology Map - Slice A



Order Details

 Order Number:
 145523120_1_1

 Customer Ref:
 10/1004

 National Grid Reference:
 335080, 389170

 Slice:
 A

 Site Area (Ha):
 0.21


 Search Buffer (m):
 1000

Site Details

Site at, Liverpool, Liverpool



Tel: Fax: Web:



Envirocheck[®] • LANDMARK INFORMATION GROUP[®]

Bedrock and Faults


Bedrock geology is a term used for the main mass of rocks forming the Earth and are present everywhere, whether exposed at the surface in outcrops or concealed beneath superficial deposits or water.

The bedrock has formed over vast lengths of geological time ranging from ancient and highly altered rocks of the Proterozoic, some 2500 million years ago, or older, up to the relatively young Pliocene, 1.8 million years ago.

The bedrock geology includes many lithologies, often classified into three types based on origin: igneous, metamorphic and sedimentary.

The BGS Faults and Rock Segments dataset includes geological faults and thin beds mapped as lines such as coal seams and mineral veins. These are not restricted by age and could relate to features of any of the 1:10,000 geology datasets.





Order Details

 Order Number:
 145523120_1_1

 Customer Ref:
 10/1004

 National Grid Reference:
 335080, 389170

 Slice:
 A

 Site Area (Ha):
 0.21


 Search Buffer (m):
 1000

Site Details

Site at, Liverpool, Liverpool



Tel: 08 Fax: 08 Web: ww



Envirocheck[®] • LANDMARK INFORMATION GROUP[®]

Combined Surface Geology

The Combined Surface Geology map combines all the previous maps into one combined geological overview of your site.


Please consult the legends to the previous maps to interpret the Combined "Surface Geology" map.

Additional Information

More information on 1:10,000 Geological mapping and explanations of rock classifications can be found on the BGS website. Using the LEX Codes in this report, further descriptions of rock types can be obtained by interrogating the 'BGS Lexicon of Named Rock Units'. This database can be accessed by following the 'Information and Data' link on the BGS website.

Contact

British Geological Survey Kingsley Dunham Centre Keyworth Nottingham NG12 5GG Telephone: 0115 936 3143 Fax: 0115 936 3276 email: enquiries@bgs.ac.uk website: www.bgs.ac.uk



Combined Geology Map - Slice A

Order Details

 Order Number:
 145523120_1_1

 Customer Ref:
 10/1004

 National Grid Reference:
 335080, 389170

 Slice:
 A

 Site Area (Ha):
 0.21

 Search Buffer (m):
 1000

Site Details

Site at, Liverpool, Liverpool



Tel: Fax: Web:

Geology 1:50,000 Maps Legends

Artificial Ground and Landslip

Map Colour	Lex Code	Rock Name	Rock Type	Min and Max Age
	WMGR	Infilled Ground	Artificial Deposit	Cenozoic - Cenozoic
\square	MGR	Made Ground (Undivided)	Artificial Deposit	Holocene - Holocene
	WGR	Worked Ground (Undivided)	Void	Holocene - Holocene

Superficial Geology

Map Colour	Lex Code	Rock Name	Rock Type	Min and Max Age
	TFD	Tidal Flat Deposits	Clay, Silty, Sandy [Unlithified Deposits Coding Scheme]	Holocene - Holocene
	TILLD	Till, Devensian	Clay, Sandy, Gravelly, Cobbly [Unlithified Deposits Coding Scheme]	Devensian - Devensian
	SSA	Shirdley Hill Sand Formation	Sand	Flandrian - Devensian

Bedrock and Faults

Map Colour	Lex Code	Rock Name	Rock Type	Min and Max Age
	TPSF Tarporley Siltsto Formation		Siltstone, Mudstone and Sandstone	Anisian - Olenekian
	HEY	Helsby Sandstone Formation	Sandstone	Anisian - Early Triassic
	СРВ	Chester Pebble Beds Formation	Sandstone, Pebbly (Gravelly)	Early Triassic - Early Triassic
	WLSF	Wilmslow Sandstone Formation	Sandstone	Early Triassic - Early Triassic
		Faults		

Envirocheck

LANDMARK INFORMATION GROUP*


Geology 1:50,000 Maps

This report contains geological map extracts taken from the BGS Digital Geological map of Great Britain at 1:50,000 scale and is designed for users carrying out preliminary site assessments who require geological maps for the area around the site. This mapping may be more up to date than previously published paper maps.

The various geological layers - artificial and landslip deposits, superficial geology and solid (bedrock) geology are displayed in separate maps, but superimposed on the final 'Combined Surface Geology' map. All map legends feature on this page. Not all layers have complete nationwide coverage, so availability of data for relevant map sheets is indicated below.

Geology 1:50,000 Maps Coverage

Map ID: Map Name: Map Name: Map Date: Bedrock Geology: Superficial Geology: Artificial Geology: Faults: Landslip: Rock Segments:	1 096 Liverpool 2006 Available Available Available Not Supplied Not Supplied		-
Geology 1:50		- Slice A	х
A16A17 A11A12		A15 A20-	×
Orde A1 A2 Customer redered		A9 A10-	
National Grid Ref Slice: Site Area (Ha): Search Buffer (m)	A 0.21	0, 389170	
Site Details: Site at, Liverpool,	Liverpool		
	ark	Tel: Fax: Web:	0844 844 9952 0844 844 9951 www.envirocheck.co.uk
v15.0 08-Nov-20	17		Page 1 of



LANDMARK INFORMATION GROUP[®]

Artificial Ground and Landslip

Artificial ground is a term used by BGS for those areas where the ground surface has been significantly modified by human activity. Information about previously developed ground is especially important, as it is often associated with potentially contaminated material, unpredictable engineering conditions and unstable ground.


Artificial ground includes:


- Made ground man-made deposits such as embankments and spoil heaps on the natural ground surface.
- Worked ground areas where the ground has been cut away such as quarries and road cuttings.
- Infilled ground areas where the ground has been cut away then wholly or partially backfilled.
- Landscaped ground areas where the surface has been reshaped.

- Disturbed ground - areas of ill-defined shallow or near surface mineral workings where it is impracticable to map made and worked ground separately.

Mass movement (landslip) deposits on BGS geological maps are primarily superficial deposits that have moved down slope under gravity to form landslips. These affect bedrock, other superficial deposits and artificial ground. The dataset also includes foundered strata, where the ground has collapsed due to subsidence.

Artificial Ground and Landslip Map - Slice A






LANDMARK INFORMATION GROUP[®]


Superficial Geology

Superficial Deposits are the youngest geological deposits formed during the most recent period of geological time, the Quaternary, which extends back about 1.8 million years from the present.

They rest on older deposits or rocks referred to as Bedrock. This dataset contains Superficial deposits that are of natural origin and 'in place'. Other superficial strata may be held in the Mass Movement dataset where they have been moved, or in the Artificial Ground dataset where they are of man-made origin.

Most of these Superficial deposits are unconsolidated sediments such as gravel, sand, silt and clay, and onshore they form relatively thin, often discontinuous patches or larger spreads.





LANDMARK INFORMATION GROUP®


Bedrock and Faults


Bedrock geology is a term used for the main mass of rocks forming the Earth and are present everywhere, whether exposed at the surface in outcrops or concealed beneath superficial deposits or water.

The bedrock has formed over vast lengths of geological time ranging from ancient and highly altered rocks of the Proterozoic, some 2500 million years ago, or older, up to the relatively young Pliccene, 1.8 million years ago.

The bedrock geology includes many lithologies, often classified into three types based on origin: igneous, metamorphic and sedimentary.

The BGS Faults and Rock Segments dataset includes geological faults (e.g. normal, thrust), and thin beds mapped as lines (e.g. coal seam, gypsum bed). Some of these are linked to other particular 1:50,000 Geology datasets, for example, coal seams are part of the bedrock sequence, most faults and mineral veins primarily affect the bedrock but cut across the strata and post date its deposition.





LANDMARK INFORMATION GROUP®

Combined Surface Geology

The Combined Surface Geology map combines all the previous maps into one combined geological overview of your site.

Please consult the legends to the previous maps to interpret the Combined "Surface Geology" map.

Additional Information

More information on 1:50,000 Geological mapping and explanations of rock classifications can be found on the BGS website. Using the LEX Codes in this report, further descriptions of rock types can be obtained by interrogating the 'BGS Lexicon of Named Rock Units'. This database can be accessed by following the 'Information and Data' link on the BGS website.

Contact

British Geological Survey Kingsley Dunham Centre Keyworth NGt12 5GG Telephone: 0115 936 3143 Fax: 0115 936 3276 email: enquiries@bgs.ac.uk website: www.bgs.ac.uk



